

DOCUMENT RESUME

ED 051 258

TM 000 296

AUTHOR Wolff, Hans  
TITLE On Stochastic Approximation.  
INSTITUTION Educational Testing Service, Princeton, N.J.  
REPORT NO RB-70-55  
PUB DATE Sep 70  
NOTE 13p.  
  
EDRS PRICE EDRS Price MF-\$0.65 HC-\$3.29  
DESCRIPTORS \*Mathematics, \*Probability, Statistical Analysis,  
\*Statistics  
IDENTIFIERS Robbins-Monro Procedure, \*Stochastic Approximation

ABSTRACT

This paper deals with a stochastic process for the approximation of the root of a regression equation. This process was first suggested by Robbins and Monro. The main result here is a necessary and sufficient condition on the iteration coefficients for convergence of the process (convergence with probability one and convergence in the quadratic mean). (Author)

ED051258

**RESEARCH**

**BULLETIN**

RB-70-55

# ON STOCHASTIC APPROXIMATION

Hans Wolff

U S DEPARTMENT OF HEALTH, EDUCATION  
& WELFARE  
OFFICE OF EDUCATION  
THIS DOCUMENT HAS BEEN REPRODUCED  
EXACTLY AS RECEIVED FROM THE PERSON OR  
ORGANIZATION ORIGINATING IT. POINTS OF  
VIEW OR OPINIONS STATED DO NOT NECESSARILY  
REPRESENT OFFICIAL OFFICE OF EDUCATION  
POSITION OR POLICY

This Bulletin is a draft for interoffice circulation. Corrections and suggestions for revision are solicited. The Bulletin should not be cited as a reference without the specific permission of the author. It is automatically superseded upon formal publication of the material.

Educational Testing Service  
Princeton, New Jersey  
September 1970

## ON STOCHASTIC APPROXIMATION

### Abstract

This paper deals with a stochastic process for the approximation of the root of a regression equation. This process was first suggested by Robbins and Monro [1].

The main result here is a necessary and sufficient condition on the iteration coefficients for convergence of the process (convergence with probability one and convergence in the quadratic mean).

## ON STOCHASTIC APPROXIMATION

### 1. Introduction and Summary

In their classical paper Robbins and Monro [1] treated the following problem.

Let  $F(y|x)$  be a family of distribution functions depending upon a real parameter  $x$ ,  $-\infty < x < +\infty$ , and let  $M(x)$ ,

$$M(x) = \int_{-\infty}^{\infty} y \, dF(y|x) \quad ,$$

be the corresponding regression function. It is assumed that  $M(x)$  and  $F(y|x)$  are unknown to the experimenter who can, however, take observations on  $F(y|x)$  for any value  $x$ . Robbins and Monro gave a method for solving stochastically the regression equation

$$(1) \quad M(x) = \alpha \quad ,$$

where  $\alpha$  is a given number. Under certain conditions on  $M(x)$  they were able to construct an iteration procedure  $\{X_n\}$  such that  $X_n$  converges in probability to the (unique) root  $\theta$  of (1).

This "Robbins-Monro procedure" is defined as follows. Let  $\{a_n\}$  be a fixed sequence of positive numbers such that

$$(2) \quad \sum_{n=1}^{\infty} a_n = \infty \quad , \quad \sum_{n=1}^{\infty} a_n^2 < \infty \quad .$$

The iteration procedure is then defined recursively as the nonstationary Markov chain  $\{X_n\}$  given by

$$(3) \quad X_{n+1} = X_n - a_n(Y_n - \alpha) \quad , \quad P(X_1 = a \in R^1) = 1 \quad ,$$

where  $Y_n$  is a random variable distributed according to  $F(y|x = X_n)$  or, in another notation,  $Y_n$  is a realization of the random variable  $Y(X_n)$ .

Later several authors (e.g., Blum [2], Dvoretzky [3]) have shown that even under weaker conditions than those imposed in [1] on  $M(x)$ , the Robbins-Monro process also converges with probability one and in the quadratic mean.

In this paper we deal with the question of whether it is possible to relax the parameter condition (2). The main result is that the condition

$$(4) \quad a_n \rightarrow 0 \quad (n \rightarrow \infty), \quad \sum_{n=1}^{\infty} a_n = \infty,$$

in connection with certain assumptions on  $M(x)$ , is necessary and sufficient for convergence with probability one and in the quadratic mean. Furthermore, the proof of convergence seems to be more elementary than proofs given by earlier writers.

## 2. Lemmas

In this section we state and prove two Lemmas which will be needed for the proof of Theorem 1 and Theorem 2 given in section 3.

Lemma A. Let  $\{a_i\}$  be a sequence of real numbers. Then

$$\sum_{i=1}^n a_i \prod_{j=i+1}^n (1 - a_j) = 1 - \prod_{i=1}^n (1 - a_i), \quad n \geq 1. \quad 1$$

Lemma B. Let  $\{a_i\}$  be a sequence of positive numbers satisfying the condition

$$a_n \rightarrow 0 \quad (n \rightarrow \infty), \quad \sum_{n=1}^{\infty} a_n = \infty.$$

Then

$$\sum_{i=1}^n a_i^2 \prod_{j=i+1}^n (1 - a_j)^2 \rightarrow 0 \quad (n \rightarrow \infty).$$

---

<sup>1</sup>Throughout this paper the factor of the last term of such a sum equals one.

Lemma A can easily be verified by induction. To prove Lemma B we note first that for any  $\varepsilon > 0$  there exists an integer  $N_0 = N_0(\varepsilon)$  such that, for all  $n \geq N_0$ ,

$$a_n < \varepsilon, \quad 0 < 1 - a_n < 1.$$

Hence we get the inequality

$$\sum_{i=1}^n a_i^2 \prod_{j=i+1}^n (1 - a_j)^2 < \sum_{i=N_0}^n (1 - a_i) \left[ \sum_{i=1}^{N_0-1} a_i^2 \prod_{j=i+1}^{N_0-1} (1 - a_j)^2 \right] + \varepsilon \sum_{i=N_0}^n a_i \prod_{j=i+1}^n (1 - a_j).$$

The factor of  $\varepsilon$  is less than one by virtue of Lemma A. Because of the divergence of  $\sum a_n$  there exists for any  $M > 0$  an integer  $N_1 = N_1(\varepsilon, M)$  such that, for all  $n \geq N_1$ ,

$$\sum_{i=N_0}^n (1 - a_i) < \varepsilon \cdot M^{-1}.$$

If we denote

$$\sum_{i=1}^{N_0-1} a_i^2 \prod_{j=i+1}^{N_0-1} (1 - a_j)^2 = M,$$

it follows immediately that

$$\sum_{i=1}^n a_i^2 \prod_{j=i+1}^n (1 - a_j)^2 < 2\varepsilon \quad \text{for all } n \geq N_1.$$

This completes the proof of Lemma B.

### 3. Stochastic Approximation of the Root of a Regression Equation

Let us assume that the regression function  $M(x)$  corresponding to the family of distribution functions  $F(y|x)$ , satisfies the following conditions:

$$(5) \quad c_1 |x - \theta| \leq |M(x) - \alpha| \leq c_2 |x - \theta| + c_3, \quad c_2 \geq c_1 > 0, \quad c_3 \geq 0;$$

$$(6) \quad \begin{aligned} M(x) &< \alpha & x < \theta \\ M(x) &= \alpha & \text{for } x = \theta \\ M(x) &> \alpha & x > \theta \end{aligned}$$

The variance of  $Y(x)$  is supposed to be uniformly bounded in  $x$ ,

$$(7) \quad \text{Var } Y(x) \leq c_4 < \infty.$$

Then we state the following theorems.

Theorem 1. If conditions (4) through (7) hold, then the stochastic process  $\{X_n\}$  given by (3) converges to  $\theta$  with probability one and in the quadratic mean,  $X_n \rightarrow \theta$  w.pr. 1,  $E(X_n - \theta)^2 \rightarrow 0$  ( $n \rightarrow \infty$ ).

If we replace condition (5) by

$$(5') \quad c_1 |x - \theta| \leq |M(x) - \alpha| \leq c_2 |x - \theta|, \quad c_2 \geq c_1 > 0,$$

and if we add the assumption that in a neighborhood of  $\theta$   $\text{Var } Y(x)$  does not vanish,

$$(8) \quad \text{Var } Y(x) \geq c_5 > 0 \quad \text{for all } x \in \{x \mid |x - \theta| < \delta, \delta > 0\},$$

then the parameter condition (4) is even necessary and sufficient for the convergence of  $\{X_n\}$  to  $\theta$ .

Theorem 2. If conditions (5'), (6), (7), (8) hold, then  $\{X_n\}$  converges to  $\theta$  with probability one and in the quadratic mean if and only if the parameter sequence  $\{a_n\}$  fulfills condition (4).

Proof of Theorem 1. We derive a recursion formula for the sequence  $E_n = E(X_n - \theta)^2$ .

From (3) we have

$$\begin{aligned} (9) \quad E_{n+1} &= E(X_{n+1} - \theta)^2 = E[X_n - \theta - a_n(Y_n - \alpha)]^2 \\ &= E_n - 2a_n E[(X_n - \theta)E(Y(x) - \alpha | x = X_n)] + a_n^2 E[E[(Y(x) - \alpha)^2 | x = X_n]] \quad . \end{aligned}$$

Because of (5) and (6) it follows that

$$\begin{aligned} E[(X_n - \theta)E(Y(x) - \alpha | x = X_n)] &= E[(X_n - \theta)(M(X_n) - \alpha)] \\ &= E[|X_n - \theta| |M(X_n) - \alpha|] \geq c_1 E(X_n - \theta)^2 \geq 0 \quad . \end{aligned}$$

From (5) and (7) we get

$$\begin{aligned} E[E[(Y(x) - \alpha)^2 | x = X_n]] &= E[E[(Y(x) - M(x) + M(x) - \alpha)^2 | x = X_n]] \\ &= E\{\text{Var } Y(X_n) + (M(X_n) - \alpha)^2\} \leq E[c_4 + c_2^2(X_n - \theta)^2 + 2c_2c_3|X_n - \theta| + c_3^2] \\ &\leq c_4 + 2c_2c_3 + c_3^2 + (c_2^2 + 2c_2c_3)E(X_n - \theta)^2 \quad . \end{aligned}$$

Using these inequalities and setting

$$c_4 + 2c_2c_3 + c_3^2 = c_6, \quad c_2^2 + 2c_2c_3 = c_7^2, \quad ,$$

it follows at once that

$$E_{n+1} \leq (1 - 2c_1a_n + c_7^2a_n^2)E_n + c_6a_n^2 \quad .$$

Because of the convergence of  $\{a_n\}$  to zero and  $c_7 \geq c_1$  there exists for each constant  $c_8$ ,  $0 < c_8 < c_1$ , an integer  $N_2$  such that for all  $n \geq N_2$

$$1 - 2c_1a_n + c_7^2a_n^2 < (1 - c_8a_n)^2 \quad .$$



This yields the more convenient inequality

$$E_{n+1} \leq (1 - c_8 a_n)^2 E_n + c_6 a_n^2, \quad n \geq N_2.$$

Adding up this inequality from  $N_2$  to  $n$  we find

$$(10) \quad E_{n+1} \leq E_{N_2} \prod_{i=N_2}^n (1 - c_8 a_i)^2 + c_6 \sum_{i=N_2}^n a_i^2 \prod_{j=i+1}^n (1 - c_8 a_j)^2.$$

The first term of the right-hand side of (10) converges to zero since  $E_{N_2}$  is finite and

$$\prod_{i=N_2}^n (1 - c_8 a_i) \rightarrow 0 \quad (n \rightarrow \infty)$$

because of the divergence of  $\sum a_i$ . Because of Lemma B the same holds for the second term,

$$\sum_{i=N_2}^n a_i^2 \prod_{j=i+1}^n (1 - c_8 a_j)^2 = (c_8)^{-2} \cdot \sum_{i=N_2}^n (c_8 a_i)^2 \prod_{j=i+1}^n (1 - c_8 a_j)^2 \rightarrow 0 \quad (n \rightarrow \infty).$$

This concludes the proof of convergence in the quadratic mean.

To show that  $\{X_n\}$  converges also with probability one we use a method which is similar to that employed by Dvoretzky [3]. We derive the convergence with probability one from the convergence in the mean.

For any pair  $\varepsilon > 0$ ,  $\delta > 0$ , there exists an integer  $N_3 = N_3(\varepsilon, \delta)$  such that, for all  $n \geq N_3$ ,

$$E_n = E(X_n - \theta)^2 < \varepsilon \delta^2.$$

We modify the sequence  $\{X_n\}$ :

$$(11) \quad X'_n = X_n \quad \text{for all } n \leq N_3$$

$$(12) \quad X'_{n+1} = \begin{cases} X'_n - a_n(Y_n - \alpha) & \text{if } |X'_n - \theta| < \delta \\ X'_n & \text{otherwise} \end{cases}, \quad n \leq N_3.$$

$Y_n$  denotes now a realization of the random variable  $Y(x = X'_n)$  instead of  $Y(x = X_n)$ .

Equations (11) and (12) imply that also

$$(13) \quad E(X'_n - \theta) < \epsilon \delta^2 \quad \text{for all } n \geq N_3.$$

If  $|X_j - \theta| \geq \delta$  for any  $j > N_3$ , it follows from (12) that  $|X'_n - \theta| \geq \delta$  for all  $n \geq j$ , and we obtain, for all  $n > N_3$ ,

$$P\left\{\max_{N_3 < j \leq n} |X_j - \theta| \geq \delta\right\} \leq P\{|X'_n - \theta| \geq \delta\}.$$

Together with (13) this implies that  $\{X_n\}$  converges with probability one to  $\theta$ , i.e.,

$$P\left\{\sup_{j > N_3} |X_j - \theta| > \delta\right\} < \epsilon.$$

This completes the proof of Theorem 1.

#### Proof of Theorem 2.

Since Theorem 1 implies the sufficiency of parameter condition 4, it remains only to prove that the parameter condition (4) is necessary for the convergence of  $\{X_n\}$  to  $\theta$ . We assume that the sequence  $\{F_n\}$  converges to zero even in the case when we use a parameter sequence  $\{a_n\}$  which does not satisfy condition (4). We show that this assumption yields a contradiction.

The parameter sequence  $\{a_n\}$  under consideration has to fulfill exactly one of the following conditions:

$$(a) \quad \sum_{i=1}^{\infty} a_i < \infty \quad ;$$

$$(b) \quad \text{there exists a subsequence } \{a_{n_i}\} \text{ and a constant } L > 0 \text{ such that}$$

$$a_{n_i} \geq L > 0 \text{ for all } i .$$

From the asserted convergence of  $E_n$  to zero it follows--as we have seen-- that  $X_n$  converges to  $\theta$  with probability one. Therefore and because of (8) there exists an integer  $N_4$  such that, with probability one,

$$\min_{n \geq N_4} \text{Var } Y(X_n) \geq c_5 > 0 .$$

In the parameter case (a), which implies  $a_n \rightarrow 0$  ( $n \rightarrow \infty$ ), we can further assume that  $N_4$  is so large that

$$0 < 1 - 2c_2 a_n + c_1^2 a_n^2 < 1 \quad \text{for all } n \geq N_4 .$$

Hence it follows from (9) by similar arguments to those used before that

$$E_{n+1} \geq E_n - 2c_2 a_n E_n + a_n^2 [E(\text{Var } Y(X_n)) + c_1^2 E_n]$$

$$\geq (1 - 2c_2 a_n + c_1^2 a_n^2) E_n + c_5 a_n^2 , \quad n \geq N_4 .$$

Again there exists for each  $c_9 > c_2$  an integer  $N_5 = N_5(c_9) \geq N_4$  such that

$$E_{n+1} \geq (1 - c_9 a_n) E_n + c_5 a_n^2 \quad \text{for all } n \geq N_5 .$$

Hence we get for the parameter case (a)

$$E_{n+1} \geq E_{N_5} \cdot \prod_{i=N_5}^n (1 - c_9 a_i)^2 + c_5 \sum_{i=N_5}^n a_i^2 \prod_{j=i+1}^n (1 - c_9 a_j)^2 \geq$$

$$c_5 \cdot f \cdot \sum_{i=N_5}^n a_i^2 = c_{10} \quad ,$$

where

$$f = \prod_{j=N_5+1}^{\infty} (1 - c_9 a_j)^2$$

is greater than zero because of the convergence of  $\sum a_i$ . Hence we have  $E_n \geq c_{10} > 0$  for all  $n \geq N_5$ , which implies the desired contradiction.

In case (r) we get the contradiction immediately by considering the sequence of inequalities

$$E_{n_i} \geq c_5 a_{n_i}^2 \geq c_5 \cdot L > 0 \quad \text{for all } n_i \geq N_4 .$$

This completes the proof of Theorem 2.

#### 4. Concluding Remarks

The crucial assumptions which lead to the weakening of the parameter condition (2) are the two assumptions contained in (5) and (5'), respectively. One of the assumptions in (5),

$$(14) \quad |M(x) - \alpha| \leq c_2 |x - \theta| + c_3 \quad ,$$

cannot be relaxed as it was pointed out, e.g., by A. Dvoretzky ([3], p. 51). However, it might be interesting to know if the validity of Theorems 1 and 2 is affected by weakening the other assumption made in (5) and (5'),

$$(15) \quad c_1 |x - \theta| \leq |M(x) - \alpha| \quad .$$

In particular, we may ask if it is possible to replace (15) by the usual condition (e.g., Blum [2], p. 382)

$$\inf_{\delta_1 \leq |x - \theta| \leq \delta_2} |M(x) - \alpha| > 0 \quad \text{for every pair of numbers } (\delta_1, \delta_2) \text{ with } 0 < \delta_1 < \delta_2 < \infty.$$

In practice, however, condition (14) and (15) will cause no trouble, because in almost all instances the experimenter knows that the root  $\theta$  lies in some finite interval  $[C_*, C^*]$ . Therefore he can replace the iteration procedure (3) by the bounded stochastic approximation process

$$X_{n+1} = \begin{cases} C_* & X_n - a_n(Y_n - \alpha) < C_* \\ X_n - a_n(Y_n - \alpha) & \text{if } C_* \leq X_n - a_n(Y_n - \alpha) \leq C^* \\ C^* & X_n - a_n(Y_n - \alpha) > C^* \end{cases}.$$

In this situation (14) and (15) do not seem very restrictive.

References

- [1] Robbins, H., and Monro, S. A stochastic approximation method. The Annals of Mathematical Statistics, 1951, 22, 400-407.
- [2] Blum, J. Approximation methods which converge with probability one. The Annals of Mathematical Statistics, 1954, 25, 382-386.
- [3] Ljvoretzky, A. On stochastic approximation. Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability I, 39-55. University of California Press, 1956.